

## 111(11) RESOLUTION UV SPECTROSCOPY FOR SOLAR SYSTEM EXPLORATION

J.M. Ajello (Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91 109; 818 354-2457; e-mail: jajello@jpluvs.jpl.nasa.gov)

I. Kanik, G. James, X. Liu, M. Ciocca, C. Noren (All at: Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91 109)  
D. Shemansky (University of Southern California, Los Angeles, CA 90089)

A new generation of high resolution UV imaging spacecraft (Polar, *Ultraviolet Imaging Telescope*) are studying the airglow and aurora of the Earth and the Jovian planets. To keep pace with these technological improvements we have developed a laboratory program to provide electron collision cross sections of the major molecular planetary gases ( $H_2$ ,  $HI$ ,  $O$ ,  $N_2$ ,  $CO_2$ ,  $SO_2$ ,  $O_2$ ,  $H_2O$  and  $CO$ ). Molecular spectra under optically thin conditions have been measured with a high resolution ( $\lambda/\Delta\lambda = 50000$ ) UV spectrometer in tandem with an electron impact collision chamber. Synthetic spectral intensities and rotational line positions for  $H_2$  based on the J-dependent transition probabilities are in good agreement with experimental intensities. A new high resolution UV model is being developed for modeling laboratory data. The model includes an accurate remeasurement of the Lyman band and Werner band cross sections. The kinetic energy distribution of  $H(2p)$  atoms resulting from electron impact dissociation of  $H_2$  has been measured. Electron impact dissociation of  $H_2$  is one of the major mechanisms leading to the observed wide profile of  $HII\alpha$  from Jupiter aurora by *Ultraviolet Imaging Telescope*. Analysis of the deconvolved line profile of  $HII\alpha$  reveals the existence of a narrow line peak (40 mÅ FWHM) and a broad pedestal base (240 mÅ FWHM). Follow-on studies of dissociative excitation of  $SHI$  259Å,  $OII$  1304Å emission from  $SO_2$  and of  $NI$  1200Å,  $NI$  1085~ from  $N_2$  also indicate substantial kinetic energy release (1 - 10 eV). We report a quantitative measurement of the predissociation fraction in the  $N_2$   $c'_4^1\Sigma_u^+$  -  $X^1\Sigma_g^+$  (0,0) band to model  $N_2$  EUV emission from Titan, Triton and the Earth. The most recent studies of the excitation cross section of the  $N_2$  ( $a^1\Pi_g$ ) state will be discussed. We report the first study of  $HII\alpha$  from atomic  $H$  in the extended energy range (10-2000 eV) where cross sections can be related to optical oscillator strengths. A small ( $\sim 50\%$ ) change in the cross section of the most abundant species at low energy ( $\sim 50$  eV) is fundamental to all electron energy loss codes.

1. 1996 Fall Meeting
2. 012102343 (J. M. Ajello)
3. (a) J.M. Ajello  
Jet Propulsion Laboratory  
MS 183-601  
Pasadena, CA 91109  
(b) Tel : 818-354-2457  
(c) Fax: 818-393-4605
4. 0
5. (a) SA03 (SPA Aeronomy)
6. Invited Talk
8. 0%
9. PO 539311
10. Invited by Session Chair  
J. Slanger for Session  
SA03: Excited atoms. . .